

## CORRIGENDUM: Integrative taxonomy resolves three new cryptic species of small southern African horseshoe bats (*Rhinolophus*)

PETER J. TAYLOR<sup>1,2\*</sup>, ANGUS MACDONALD<sup>2,6</sup>, STEVEN M. GOODMAN<sup>3,4</sup>,  
TERESA KEARNEY<sup>5,6</sup>, FENTON P. D. COTTERILL<sup>7</sup>, SAM STOFFBERG<sup>8</sup>,  
ARA MONADJEM<sup>9,10</sup>, M. CORRIE SCHOEMAN<sup>2</sup>, JENNIFER GUYTON<sup>11</sup>,  
PIOTR NASKRECKI<sup>12</sup>, ADRIAAN ENGELBRECHT<sup>13</sup> and LEIGH R. RICHARDS<sup>14</sup>

<sup>1</sup>School of Mathematical & Natural Sciences, University of Venda, Private Bag X5050, Thohoyandou 0950, South Africa

<sup>2</sup>School of Life Sciences, University of KwaZulu-Natal, Biological Sciences Building, South Ring Road, Westville Campus, Durban, KwaZulu-Natal 3630, South Africa

<sup>3</sup>Field Museum of Natural History, 1400 South Lake Shore Drive, Chicago, IL 60605, USA

<sup>4</sup>Association Vahatra, BP 3972, Antananarivo 101, Madagascar

<sup>5</sup>Ditsong National Museum of Natural History, PO Box 413, Pretoria 0001, South Africa

<sup>6</sup>School of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Private Bag 3, Wits 2050, South Africa

<sup>7</sup>Geocodynamics Research Hub, c/o Department of Botany and Zoology, University of Stellenbosch, Private Bag X1 Matieland, 7602, Stellenbosch, South Africa

<sup>8</sup>Department of Botany and Zoology, University of Stellenbosch, Private Bag X1 Matieland, 7602, Stellenbosch, South Africa

<sup>9</sup>Department of Biological Sciences, University of Eswatini, Private Bag 4, Kwaluseni, Eswatini

<sup>10</sup>Mammal Research Institute, Department of Zoology & Entomology, University of Pretoria, Private Bag 20, Hatfield 0028, Pretoria, South Africa

<sup>11</sup>Department of Ecology and Evolutionary Biology, Princeton University, 106A Guyot Hall, Princeton, NJ 08544, USA

<sup>12</sup>Museum of Comparative Zoology, Harvard University, 26 Oxford Street, Cambridge, MA 02138, USA

<sup>13</sup>University of Western Cape, New Science Building, Core 2 Level 4, Robert Sobukwe Road, Cape Town, 7535, South Africa


<sup>14</sup>Durban Natural Science Museum, PO Box 4085, Durban, 4001, South Africa

This corrigendum advises that the cytochrome *b* sequence, GenBank accession number MG980682, of *Rhinolophus landeri* Martin, 1837 from Liberia (accession number DM12593 in the Durban Natural Science Museum), reported by Taylor *et al.* (2018) is contaminated and represents a sequence of *Bos taurus* Linnaeus, 1758 rather than the horseshoe bat *R. landeri*. In this corrigendum, we re-analysed the molecular data from Taylor *et al.* (2018), replacing the contaminated sequence with a sequence of *R. landeri* from Mali in West Africa (GenBank accession number

KU531353), which truly represents *R. landeri* (see Dool *et al.*, 2016).

The revised data do not change the original taxonomic conclusions of Taylor *et al.* (2018). They show that *R. landeri* s.s. is the sister species (although not strongly supported by maximum likelihood or parsimony bootstrap values and moderately supported by Bayesian probability) of newly described *Rhinolophus lobatus* from southern Africa (Fig. 1, corresponding to the corrected version of Fig. 2 of Taylor *et al.*, 2018). A similar close relationship was also shown by Dool *et al.* (2016), who compared western and southern African populations of *R. landeri* s.l. (*R. landeri* and *R. lobatus* Peters, 1852, respectively, in our interpretation).

\*Corresponding author. E-mail: [peter.taylor@univen.ac.za](mailto:peter.taylor@univen.ac.za)



**Figure 1.** Bayesian tree of partial *Cytb* sequences, with nodal support values based on Bayesian probabilities (in bold to the right of nodes) and bootstrap values for maximum likelihood (left of node and above branch) and maximum parsimony (left of node and below branch).

**Table 1.** Mean uncorrected *p*-distances (below) diagonal and Kimura 2-parameter model-corrected distances (above diagonal) based on 667 cytochrome *b* (*Cytb*) sequences between eight recognized and newly proposed species of small African *Rhinolophus* bats

| <i>Rhinolophus</i> | <i>rhodesiae</i> | <i>simulator</i> | <i>denti</i>  | <i>swinnnyi</i> | <i>capensis</i> | <i>lobatus</i> | <i>gorongosae</i> | <i>landeri</i> |
|--------------------|------------------|------------------|---------------|-----------------|-----------------|----------------|-------------------|----------------|
| <i>rhodesiae</i>   | <b>0.0024</b>    | 0.003            | 0.050         | 0.092           | 0.088           | 0.156          | 0.075             | 0.151          |
| <i>simulator</i>   | 0.003            | <b>0.0012</b>    | 0.049         | 0.093           | 0.087           | 0.156          | 0.072             | 0.148          |
| <i>denti</i>       | 0.048            | 0.047            | <b>0.0071</b> | 0.072           | 0.068           | 0.138          | 0.107             | 0.130          |
| <i>swinnnyi</i>    | 0.085            | 0.085            | 0.067         | <b>0.0110</b>   | 0.060           | 0.143          | 0.145             | 0.147          |
| <i>capensis</i>    | 0.081            | 0.080            | 0.064         | 0.057           | <b>0.0056</b>   | 0.149          | 0.135             | 0.148          |
| <i>lobatus</i>     | 0.137            | 0.137            | 0.124         | 0.128           | 0.132           | <b>0.0028</b>  | 0.194             | 0.055          |
| <i>gorongosae</i>  | 0.071            | 0.069            | 0.099         | 0.130           | 0.123           | 0.168          | <b>0.0179</b>     | 0.176          |
| <i>landeri</i>     | 0.133            | 0.131            | 0.117         | 0.131           | 0.131           | 0.052          | 0.155             | –              |

Within-species uncorrected *p*-distances are provided in bold face on the diagonals, except for *R. landeri* where only one individual was available.

The two sister species show a *p*-divergence of 5.3%, similar to that found between other species pairs of well-recognized species, e.g. *Rhinolophus capensis* Lichtenstein, 1823 and *Rhinolophus swinnnyi* Gough, 1908 (*p* = 5.7%) and *Rhinolophus denti* Thomas, 1904 and *Rhinolophus rhodesiae* Roberts, 1946 (*p* = 4.8%) (Table 1, corresponding to corrected version of Table 1 of Taylor *et al.*, 2018). Together with morphological differences described in the original paper, this divergence supports the recognition of *R. lobatus* as a species distinct from *R. landeri*.

#### ACKNOWLEDGEMENTS

We thank Bruce Patterson and Terry Demos for picking up and sharing the sequencing error. PJT acknowledges the financial support of the University of Venda, the South African National Research Foundation and the Department of Science and Technology under the South African Research

Chair Initiative (SARChI) on Biodiversity Value and Change within the Vhembe Biosphere Reserve hosted at University of Venda and co-hosted by the Centre for Invasion Biology at University of Stellenbosch.

#### REFERENCES

- Dool SE, Puechmaile SJ, Foley NM, Allegrini B, Bastian A, Mutumi GL, Maluleke TG, Odendaal LJ, Teeling EC, Jacobs DS. 2016. Nuclear introns outperform mitochondrial DNA in inter-specific phylogenetic reconstruction: lessons from horseshoe bats (Rhinolophidae: Chiroptera). *Molecular Phylogenetics and Evolution* **97**: 196–212.
- Taylor PJ, Macdonald A, Goodman SM, Kearney T, Cotterill FP, Stoffberg S, Monadjem A, Schoeman MC, Guyton J, Naskrecki P, Richards LR. 2018. Integrative taxonomy resolves three new cryptic species of small southern African horseshoe bats (*Rhinolophus*). *Zoological Journal of the Linnean Society* **184**: 1249–1276.